

# BANGLADESH JOURNAL OF NEUROSURGERY

□ VOLUME 5 □ NUMBER 2 □ JANUARY 2016

## CONTENTS

|   |    |
|---|----|
| <b>Editorial</b>  | 29 |
| <b>Original Articles</b>  |    |
| Primary bone tumour of the spine: Surgical treatment and Histopathological analysis<br><i>Md. Shafiul Alam, Md. Shafiqul Kabir Khan, Monsur Ahmed, Muhammad Shamsul Arefin, Kazi Hafiz Uddin, Mohammad Hossain, Md. Rezaul Amin</i>   | 31 |
| Outcome of Conservative Management of Traumatic Acute Subdural Haematoma in Relation to Underlying Brain Injury<br><i>Sanat Kumar saha, Mohammad Abu Sayed, Joynul Islam, Shafiul Alam, ATM Asadullah, Misbah Uddin Ahamed, Shamsul Islam Khan</i>                                  | 37 |
| Signs and Symptoms of Early VP Shunt Infection in a Tertiary Hospital<br><i>Kaisar Haroon, Tania Taher, Mokhlasur Rahman, Kanak Kanti Barua</i>   | 41 |
| Assessment of Serum Homocysteine Level in Epileptic Patients on Carbamazepine Treatment<br><i>Md. Abu Hanif, M A Hannan, M Iqbal Arslan, SK Mahbub Alam, Mohammad Hossain, Md. Aynul Hoque, Niloy Ranjan Roy</i>  | 44 |
| <b>Case Reports</b>   |    |
| Tubercular Spinal Epidural Abscess Involving the Lumbo-Sacral Region: A Case Report with Review of Literature<br><i>Akhalaque Hossain Khan, Ranjit Kumar Chaurasiya, Narendra Shalike, Mohammed Aminul Hasnat, Bipin Chaurasia, Abhishek Chaturbedi, Asifur Rahman, Rezaul Amin</i> | 48 |
| Giant Ganglioneuroma of Thoracic Spine: A Case Report<br><i>Das S, Alam K, , Rashid MM, Islam MM, Mahbub H, Khan MSI</i>  | 51 |



**OFFICIAL ORGAN OF  
BANGLADESH SOCIETY OF NEUROSURGEONS**

# Primary Bone Tumour of the Spine: Surgical Treatment and Histopathological Analysis

Md. Shafiul Alam<sup>1</sup>, Md. Shafiqul Kabir Khan<sup>2</sup>, Monsur Ahmed<sup>3</sup>, Muhammad Shamsul Arefin<sup>4</sup>, Kazi Hafiz Uddin<sup>5</sup>, Mohammad Hossain<sup>6</sup>, Md. Rezaul Amin<sup>6</sup>

### Abstract

**Background:** Primary bone tumour of the spine are rare but not uncommon. These include both benign and malignant lesions. Surgical resection and stabilization in some cases are the treatment of choice. Adjuvant radio and/or chemotherapy are needed in malignant cases.

**Methods:** We retrospectively reviewed all the primary bone tumour of the spine that were surgically treated in the department of Neurosurgery of Dhaka Medical College Hospital and National Institute of Neurosciences and Hospital Dhaka, Bangladesh between 2010 and 2015.

**Results:** Eighteen cases were included that presented at a median age of 36 years (range 14 to 49 years). Pain was the most common presenting symptom. Three patients had significant neurological deficits at time of presentation and in most cases there was an improvement after surgery. A variety of surgical strategies was employed with the use of adjuvant radio- or chemotherapy in six cases. Six benign and twelve malignant tumours were treated.

**Conclusions:** Primary bone tumours of the spine are often associated with a significant delay in diagnosis. Surgical strategy should be individualized for each case. Acceptable results can be achieved with early surgery and adequate chemo-radiotherapy.

**Key words:** primary bone tumour, spine, surgical treatment, histopathology.

*Bang. J Neurosurgery 2016; 5(2): 31- 36*

### Introduction:

Primary bone tumour of the spine is a heterogenous group of conditions that includes both benign and malignant tumours, with only 6% of these tumours considered malignant.<sup>1</sup> The spine is a frequent site of metastasis, but primary bone tumours of the spine are rare and make up only 4.2% of spinal tumours.<sup>2</sup> Eleven per cent (11%) of primary musculoskeletal tumours occur in the spine with certain tumours having an affinity for the spine. Although rare, these tumours offer a considerable diagnostic dilemma and therapeutic challenge. Timely diagnosis is crucial, as immediate attention could affect overall prognosis.<sup>3</sup>

The most common primary tumour of the spine is metastatic deposits. A number of both benign and

malignant tumours may arise primarily from the spine. Among benign tumour osteoid osteoma, osteoblastoma, osteochondroma, giant cell tumour (GCT), aneurysmal bone cyst (ABC), eosinophilic granuloma (EG), haemangioma are common. Among malignant lesion malignant lymphoma, sarcoma, ewing sarcoma, osteosarcoma, chondrosarcoma, malignant fibrous histiocytoma (MFH), fibrosarcoma, chordoma, plasmacytoma/multiple myeloma and peripheral primitive neuroectodermal tumour (pPNET) are common.

Primary bone tumours are more common in the thoracic and lumbosacral regions than in the cervical spine.<sup>4</sup> These tumours occur according to a typical anatomical distribution within the vertebra. In general, malignant tumours occur more frequently in the anterior elements and benign tumours in the posterior elements.<sup>3-5</sup>

The aims of treatment of these axial neoplasms include complete resection if feasible, preservation of neurological function and stabilisation of the spine if needed.<sup>4</sup> The aim of our study was to review the primary bone tumours of the spine treated at our unit regarding the presentation, the surgical strategy employed and the outcome.

### Materials and Methods:

This is a retrospective study. All the primary bone tumours of the spine that received surgical treatment

1. Associate Professor, Department of Gamma Knife, National Institute of Neurosciences & Hospital.

2. Associate Professor, Department of Neurosurgery, Dhaka Medical College Hospital.

3. Resident Surgeon, Department of Neurosurgery, National Institute of Neurosciences & Hospital.

4. Junior Consultant, Department of Neuroanaesthesiology, National Institute of Neurosciences & Hospital.

5. Registrar, Department of Clinical Neurosurgery, National Institute of Neurosciences & Hospital.

6. Associate Professor, Department of Neurosurgery, Bangabandhu Sheikh Mujib Medical University, Dhaka

**Address of Correspondence:** Associate Professor, Department of Gamma Knife, National Institute of Neurosciences & Hospital, E mail: dr\_chapal@hotmail.com, Tel: 01711567358.

at the department of Neurosurgery of Dhaka Medical College Hospital and National Institute of Neurosciences and Hospital Dhaka, Bangladesh from 2010 to 2015 were included in this study. We excluded all secondary deposition in the spine. We treated 18 cases during this period that were included according to these criteria. A case note and imaging review was done where patient demographics, details of presentation, clinical information and the management strategy were analyzed. All complications and mortalities were recorded.

### Results:

Between 2010 and 2015, a total 18 patients were diagnosed as primary bone tumour of the spine. The median age at presentation was 36 years (range 14 to 49 years) with eleven(61.12%) male and seven(38.88%) female patients affected. The median follow-up was 18 months (range 3 to 60 months).

**Table-I**  
*Distribution of patient by age.*

| Age group      | Frequency | Percentage |
|----------------|-----------|------------|
| 0 to 10 years  | 0         | 0          |
| 11 to 20 years | 3         | 16.67      |
| 21 to 30 years | 7         | 38.89      |
| 31 to 40 years | 4         | 22.22      |
| 41 to 50 years | 4         | 22.22      |
| Total          | 18        | 100        |

**Table-II**  
*Distribution of patient by sex.*

| Sex    | Frequency | Percentage |
|--------|-----------|------------|
| Male   | 11        | 61.12      |
| Female | 7         | 38.88      |
| Total  | 18        | 100        |

**Table-III**  
*Distribution of patient by site of involvement.*

| Site of involvement | Frequency | Percentage |
|---------------------|-----------|------------|
| Cervical            | 1         | 5.55       |
| Dorsal              | 7         | 38.89      |
| Lumber              | 6         | 33.34      |
| Sacrum              | 4         | 22.22      |
| Total               | 18        | 100        |

The most utilized diagnostic tool was MRI and CT scan of the spine. It helped to diagnose the case as well as planning of surgery. X ray is no longer a standard diagnostic tool of spine tumour, it was used in all cases as complimentary diagnostic tool. It also helped to find out the extra spinal involvement like rib cases in two cases. Pre-operative CT guided FNAC was done in seven cases. Post-operative histopathological study was done in all cases.

Primary bone tumour of the spine most commonly involved the dorsal (38.89%) and the Lumber (33.34%). It affect the sacral (22.22%) and the cervical spine (5.55%) less frequently. (Table-3)


**Table-IV**  
*Distribution of patient by histopathological findings.*

| Type      | Histopathology          | Frequency | percentage |
|-----------|-------------------------|-----------|------------|
| Benign    | -Osteoid osteoma        | 1         | 5.55       |
|           | -Haemangioma            | 1         | 5.55       |
|           | -Giant cell tumour      | 1         | 5.55       |
|           | -Aneurysmal bone cyst   | 2         | 11.11      |
|           | -Eosinophilic granuloma | 1         | 5.55       |
| Malignant | -Myxoid Chondrosarcoma  | 1         | 5.55       |
|           | -ewing's sarcoma        | 2         | 11.11      |
|           | -Chondrosarcoma         | 2         | 22.22      |
|           | -Chordoma               | 3         | 33.33      |
|           | -Multiple myeloma       | 2         | 22.22      |
|           | -pNET                   | 2         | 22.22      |
| Total     |                         | 18        | 100        |


Pain was the most common presenting symptom (15 from 18 cases) and four cases presented with pain alone. Pain was mostly axial pain in the affected region (11 of 18 cases), although three patients presented with chest pain and one with flank pain. Other presenting symptoms included neurological symptoms and odynophagia. One patient presented with a recurrence of a malignant aggressive tumour (Ewing sarcoma of sacral spine) following incomplete resection at another unit. Ten patients had no neurological deficit at time of presentation. Two patients presented with radicular symptoms and another three had significant neurological deficits at time of presentation. One case presented with significant neurological deficit with a diagnosis of benign aggressive lesions (Aneurysmal bone cyst). Two of these cases had significant improvement after surgery. We had no cases of neurological deterioration following surgery. We treated six benign and twelve malignant tumours. The histopathological diagnoses are summarised in Table 4. In all cases the surgery was performed by the senior author. A

variety of surgical strategies was utilised. An intralesional resection or debulking of the tumour was performed in six cases and a marginal resection was performed in nine cases to correlate with diagnosis. In six patients a fusion was performed by transpedicular screws and rods. Adjuvant therapy was used where indicated. Two of the recurrent cases


were in malignant tumours and one in benign aggressive tumour. In the one recurrent malignant tumour a repeat attempted marginal resection was performed. The other two recurrent cases were not re-operated as resection was not possible with acceptable morbidity. We had three deaths in the series on further follow up.




**Fig.-1: CT Scan of the lumbosacral spine showing Sacral Chordoma.**




**Fig.-2: MRI of the dorsal spine showing Ewing Sarcoma of D-9 vertebrae with involvement of ribs.**



**Fig.-3:** MRI showing extensive Chondrosarcoma of the dorsal spine.



**Fig.-4:** MRI of the dorsolumbar spine showing Aneurysmal Bone Cyst of the L-2 vertebrae.



**Fig.-5:** Pre and post-Operative X-ray of L-2 Aneurysmal Bone Cyst.

**Discussion:**

The most common presenting complaint of patients with primary bone tumour of the spine is pain, with roughly 60% complaining of axial pain and 25% of radicular symptoms.<sup>3</sup> The presence of a neurological deficit carries a worse prognosis.<sup>11</sup> Pain at rest or night pain is red flag symptoms that warrant further investigation. Spinal deformity is obvious when present, but occurs in less than 10% of patients. The most common cause of a painful scoliosis in adolescents is an osteoid osteoma.<sup>5</sup> A delay in diagnosis is common and our finding is confirmed in other series.<sup>4</sup> This is due to the presenting symptoms often being vague, pain often being the only symptom, and the initial radiographs often only showing subtle or no abnormalities leading to the symptoms not being properly investigated. Most patients report a slow, gradual onset of pain in the involved area. In benign tumours of the spine an average of 26 weeks of symptoms before presentation are reported.<sup>4</sup> This was confirmed in our series where the average delay in presentation was 7 months.

A variety of classifications systems is used for primary bone tumours of the spine. The Enneking staging system<sup>12</sup> was initially described for primary bone tumours of the appendicular skeleton, but is applied to tumours of the spine as well. It prognosticates as well as guides surgical resection margins and has been validated in the literature. The Weinstein-Boriani-Biagini surgical staging system<sup>13</sup> is an anatomical classification based on the Enneking system that describes the number of vertebra as well as the number of sectors within a vertebra involved. Recommendations are then made for the most appropriate approach to resection of the lesion. We used the Enneking staging system. According to this system we treated six benign and twelve malignant tumours. Of the benign group five could be classified as active and seven as aggressive. Appropriate treatment may be observational (Enneking grade 0 latent lesions) or surgical (most other lesions), depending on the level of pain, instability, neurologic compromise and the natural history of the lesion.<sup>5</sup> The aims of surgery are complete resection of the lesion where feasible with preservation of neurological function. We used a variety of surgical strategies customised to the patient and the tumour. In peripheral malignant sarcomas the most important factor affecting survival is complete resection of the tumour with a wide

margin.<sup>14,15</sup> There are however no reports prescribing surgical margins in spinal tumours. Traditional intralesional resection in a piecemeal fashion is thought to increase the likelihood of local recurrence. True wide resection would cause unacceptable morbidity as it would involve resecting the segment of spinal cord. Total en bloc spondylectomy in a single stage from an all posterior approach was described by Tomita.<sup>14</sup> Here wide margins are achieved except at the pedicles and occasionally the spinal canal. This approach decreases local recurrence rates.<sup>16-18</sup> In benign aggressive lesions surgical eradication also provides the best long-term cure.<sup>6,18</sup> We had three cases of local recurrence in our series. Two were in malignant tumours where a marginal resection was performed. The other recurrent case was a thoracic giant cell tumour encapsulating the aorta and forcing the surgeon to perform an intralesional resection.

Recent advances in chemotherapy have led to improved survival in malignant primary bone tumours, including in the spine.<sup>7,19</sup> Adjuvant radiotherapy is indicated in some malignant and benign aggressive lesions.<sup>6,9</sup> The use of local radiotherapy does increase the risk of major local complications, including sepsis.<sup>8,10</sup> We used radio- and chemotherapy in a number of cases as guided by our local oncology unit. The main limitations of this study are the retrospective character and the relatively small numbers.

**Conclusion:**

Most primary bone tumour of the spine in our population occurred in adults and was malignant. Surgical treatment resulted in pain reduction and neurological improvement, allowing a rapid adjuvant therapy in good clinical condition. An experienced multidisciplinary team of tertiary care center should perform diagnosis and treatment of these extremely rare tumors. In order to optimize management and elaborate guidelines, we encourage national or international registries, so that correct epidemiology, better classification and understanding, improved diagnostics, superior treatment and an overall better outcome may be achieved. We conclude that primary bone tumours of the spine are often associated with a significant delay in diagnosis. Surgical strategy and adjuvant therapy should be individualized for each case. Acceptable results with minimal complications can be achieved with this approach.

**References:**

1. Boriani S, Biagini R, De Iure F, et al. Primary bone tumors of the spine: a survey of the evaluation and treatment at the Istituto Ortopedico Rizzoli. *Orthopedics*. 1995;18(10):993–1000.
2. Davis JH. Anatomical classification and surgical considerations: Primary spinal tumours. An overview. *South African Med J*. 2011;10(3):26–30.
3. Sansur CA, Pouratian N, Dumont AS, Schiff D, Shaffrey CI, Shaffrey ME. Part II: spinal-cord neoplasms—primary tumours of the bony spine and adjacent soft tissues. *Lancet Oncol*. 2007;8(2):137–47.
4. Beer SJ, Menezes AH. Primary tumors of the spine in children: Natural history, management and long-term follow-up. *Spine (Phila Pa 1976)*. 1997;22(6):649–58.
5. Thakur N a, Daniels AH, Schiller J, et al. Benign tumors of the spine. *J Am Acad Orthop Surg*. 2012;20(11):715–24.
6. Harrop JS, Schmidt MH, Boriani S, Shaffrey CI. Aggressive ‘benign’ primary spine neoplasms. Osteoblastoma, aneurysmal bone cyst and giant cell tumour. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S39–S47.
7. Fisher CG, Andersson GBJ, Weinstein JN. Spine focus issue. Summary of management recommendations in spine oncology. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S2–S6.
8. Fisher CG, Keynan O, Ondra S, Gokaslan Z. Introduction to focus issue in spine oncology: the synthesis of evidence and expert opinion for best practice recommendation. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S21–S25.
9. Choi D, Crockard A, Bunger C, et al. Review of metastatic spine tumour classification and indications for surgery: the consensus statement of the Global Spine Tumour Study Group. *Eur Spine J*. 2010;19(2):215–22.
10. Boriani S, Bandiera S, Donthineni R, et al. Morbidity of en bloc resections in the spine. *Eur Spine J*. 2010;19(2): 231–41.
11. Kelley SP, Ashford RU, Rao AS, Dickson RA. Primary bone tumours of the spine: a 42-year survey from the Leeds Regional Bone Tumour Registry. *Eur Spine J*. 2007;16(3):405–409.
12. Enneking WF, Spanier SS, Goodman MA. A system for the surgical staging of musculoskeletal sarcoma. *Clin Orthop Relat Res*. 1980;153:106–20.
13. Boriani S, Weinstein JN, Biagini R. Primary bone tumors of the spine: Terminology and surgical staging. *Spine (Phila Pa 1976)*. 1997;22(9):1036–44.
14. Tomita K, Kawahara N, Baba H, Tsuchiya H, Fujita T, Toribatake Y. Total en bloc spondylectomy: a new surgical technique for primary malignant tumors. *Spine (Phila Pa 1976)*. 1997;22(3):324–33.
5. Gore L, Greffe BS, Rothenberg SS, Erickson M, Schreiber DP, Handler MH. Long-term survival after intralesional resection and multi-modal therapy of thoracic spine osteosarcoma. *Med Pediatr Oncol*. 2003; (6):400–402.
16. Boriani S, Saravanja D, Yamada Y, Varga PP, Biagini R, Fisher CG. Challenges of local recurrence and cure in low grade malignant tumors of the spine. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S48–S57.
17. Sciubba DM, Okuno SH, Dekutoski MB, Gokaslan ZL. Ewing and osteogenic sarcoma: evidence for multidisciplinary management. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S58–S68.
18. Yamazaki T, McLoughlin GS, Patel S, Rhines LD, Fournay DR. Feasibility and safety of en bloc resection for primary spine tumors: a systematic review by the Spine Oncology Study Group. *Spine (Phila Pa 1976)*. 2009;34(22 Suppl):S31–S38.
19. Kim HJ, McLawhorn AS, Goldstein MJ, Boland PJ. Malignant osseous tumors of the pediatric spine. *J Am Acad Orthop Surg*. 2012;20:646–56. 20. Bremnes RM, Hauge HN, Sagsvee R. Radiotherapy in the Treatment of Symptomatic Vertebral Hemangiomas: Technical Case Report. *Neurosurgery*. 1996;39(5):1054–58.